

# Trig. Past Papers Unit 2 Outcome 3

## Written Questions

- [SQA] 1. Solve the equation  $3 \cos 2x^\circ + \cos x^\circ = -1$  in the interval  $0 \leq x \leq 360$ .

5

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
	5	A/B	CR	T10	60, 131·8, 228·2, 300	2000 P2 Q5

<ul style="list-style-type: none"> <li>•<sup>1</sup> ss: know to use <math>\cos 2x = 2\cos^2 x - 1</math></li> <li>•<sup>2</sup> pd: process</li> <li>•<sup>3</sup> ss: know to/and factorise quadratic</li> <li>•<sup>4</sup> pd: process</li> <li>•<sup>5</sup> pd: process</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>3(2\cos^2 x^\circ - 1) + \cos x^\circ = -1</math></li> <li>•<sup>2</sup> <math>6\cos^2 x^\circ + \cos x^\circ - 2 = 0</math></li> <li>•<sup>3</sup> <math>(2\cos x^\circ - 1)(3\cos x^\circ + 2)</math></li> <li>•<sup>4</sup> <math>\cos x^\circ = \frac{1}{2}, x = 60, 30</math></li> <li>•<sup>5</sup> <math>\cos x^\circ = -\frac{2}{3}, x = 132, 228</math></li> </ul>
--	--

- [SQA] 2. Solve the equation  $\cos 2x^\circ + 5 \cos x^\circ - 2 = 0$ ,  $0 \leq x < 360$ .

5

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source  1994 P1 qu.15
		C	A/B	C	A/B	C	A/B		
5	2.3			1	4			2.3.5	

<ul style="list-style-type: none"> <li>•<sup>1</sup> Replacing <math>\cos 2x</math> by <math>2\cos^2 x - 1</math></li> <li>•<sup>2</sup> <math>2\cos^2 x + 5\cos x - 3 = 0</math></li> <li>•<sup>3</sup> <math>(2\cos x - 1)(\cos x + 3) = 0</math></li> <li>•<sup>4</sup> <math>60^\circ</math></li> </ul>	<ul style="list-style-type: none"> <li>•<sup>5</sup> <math>300^\circ</math> and no extraneous solutions and no solution for <math>\cos x = -3</math> indicated. (If a reason is given, it must be valid).</li> </ul>
---	--

- [SQA] 3. Find the exact solutions of the equation  $4 \sin^2 x = 1$ ,  $0 \leq x < 2\pi$ .

4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source  1995 P1 qu.8
		C	A/B	C	A/B	C	A/B		
4	2.3	4						2.3.1      1.2.1	

<ul style="list-style-type: none"> <li>•<sup>1</sup> know to factorise, take square roots</li> <li>•<sup>2</sup> <math>\sin x = \frac{1}{2}</math></li> <li>•<sup>3</sup> <math>x = \frac{\pi}{6}, \frac{5\pi}{6}</math></li> <li>•<sup>4</sup> <math>\sin x = -\frac{1}{2}</math> and <math>x = \frac{7\pi}{6}, \frac{11\pi}{6}</math></li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> replace <math>\sin^2 x</math> by <math>\frac{1}{2}(1 - \cos 2x)</math></li> <li>•<sup>2</sup> <math>\cos 2x = \frac{1}{2}</math></li> <li>•<sup>3</sup> <math>2x = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \frac{11\pi}{3}</math></li> <li>•<sup>4</sup> <math>x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}</math></li> </ul>
--	---

[SQA] 4. Solve the equation  $2\cos^2 x = \frac{1}{2}$ , for  $0 \leq x \leq \pi$ .

3

part marks	Unit	non-calc		calc		calc neut		Content Reference :		2.3
		C	A/B	C	A/B	C	A/B	Main	Additional	
.	3	2.3	3					2.3.1	1.2.1	Source 1990 P1 qu.15
$\bullet^1 \cos x = \pm \frac{1}{2}$ $\bullet^2 x = \frac{\pi}{3}$ $\bullet^3 \frac{2\pi}{3}$										

[SQA] 5. Solve the equation  $\cos 2x^\circ + \cos x^\circ = 0$ ,  $0 \leq x < 360$ .

5

part marks	Unit	non-calc		calc		calc neut		Content Reference :		2.3
		C	A/B	C	A/B	C	A/B	Main	Additional	
.	5	2.3				5		2.3.5		Source 1995 P1 qu.15
$\bullet^1$ substitute $2\cos^2 x^\circ - 1$ for $\cos 2x^\circ$ $\bullet^2 (2\cos x^\circ - 1)(\cos x^\circ + 1) = 0$ $\bullet^3 \cos x^\circ = \frac{1}{2}, \cos x^\circ = -1$ $\bullet^4 x = 60, 300$ $\bullet^5 x = 180$										

[SQA] 6. Solve  $2\sin 3x^\circ - 1 = 0$  for  $0 \leq x \leq 180$ .

4

part marks	Unit	non-calc		calc		calc neut		Content Reference :		2.3
		C	A/B	C	A/B	C	A/B	Main	Additional	
.	4	2.3	4					2.3.1	1.2.1	Source 1989 P1 qu.7
$\bullet^1 \sin 3x^\circ = 0.5$ $\bullet^2 3x = 30, 150$ $\bullet^3 x = 10, 50$ $\bullet^4$ solution is 10, 50, 130										

[SQA] 7.

(a) Show that  $2 \cos 2x^\circ - \cos^2 x^\circ = 1 - 3 \sin^2 x^\circ$ .

2

(b) Hence solve the equation  $2 \cos 2x^\circ - \cos^2 x^\circ = 2 \sin x^\circ$  in the interval  $0 \leq x < 360$ .

4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
(a) 2	2.3			1	1			2.3.3	
(b) 4	2.3			1	3			2.3.5	Source 1997 P1 qu.18

- <sup>1</sup> substitute  $1 - 2 \sin^2 x^\circ$  for  $\cos 2x^\circ$
- <sup>2</sup> substitute  $1 - \sin^2 x^\circ$  for  $\cos^2 x^\circ$
- <sup>3</sup>  $3 \sin^2 x^\circ + 2 \sin x^\circ - 1 = 0$
- <sup>4</sup>  $(3 \sin x^\circ - 1)(\sin x^\circ + 1) = 0$
- <sup>5</sup>  $\sin x^\circ = \frac{1}{3}, -1$
- <sup>6</sup>  $19.5^\circ, 160.5^\circ, 270^\circ$

[SQA] 8. Solve the equation  $\sin 2x^\circ + \sin x^\circ = 0$ ,  $0 \leq x < 360$ .

5

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
. 5	2.3	5						2.3.5	Source 1996 P1 qu.10

- <sup>1</sup>  $2 \sin x \cos x + \sin x = 0$
- <sup>2</sup>  $\sin x(2 \cos x + 1) = 0$
- <sup>3</sup>  $\sin x = 0, \cos x = -\frac{1}{2}$
- <sup>4</sup> 1st:  $x = 0, 180$
- <sup>5</sup> 2nd:  $x = 120, 240$

[SQA] 9. Find, correct to one decimal place, the value of  $x$  between 180 and 270 which satisfies the equation  $3 \cos(2x^\circ - 40^\circ) - 1 = 0$ .

5

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
. 5	2.3			5				2.3.1	Source 1992 P1 qu.5

- <sup>1</sup>  $\cos(2x - 40)^\circ = \frac{1}{3}$
- <sup>2</sup>  $\cos^{-1} \frac{1}{3} = 70.53$
- <sup>3</sup>  $2x - 40 = 70.5^\circ, 289.5^\circ, 430.5^\circ, 649.5^\circ$
- <sup>4</sup>  $x = 55.25^\circ, 164.75^\circ, 235.25^\circ, 344.75^\circ$
- <sup>5</sup>  $x = 235.25^\circ$

[SQA] 10.

- (a) Write the equation  $\cos 2\theta + 8 \cos \theta + 9 = 0$  in terms of  $\cos \theta$  and show that, for  $\cos \theta$ , it has equal roots. 3

- (b) Show that there are no real roots for  $\theta$ . 1

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional		2.3 Source 1998 P1 qu.18
		C	A/B	C	A/B	C	A/B	2.3.3	2.1.6	
(a) 3	2.3					1	2	2.3.3	2.1.6	
(b) 1	1.2					1	1	1.2.1		

$\bullet^1 2\cos^2 \theta - 1 + 8\cos \theta + 9$   
 $\bullet^2 2(\cos \theta + 2)^2 = 0$   
 or "  $b^2 - 4ac = 16 - 4 \times 1 \times 4$   
 $\bullet^3 \cos \theta = -2$  twice or "  $b^2 - 4ac = 0$   
 $\bullet^4 \cos \theta = -2$  has no solution

[SQA] 11. If  $f(a) = 6 \sin^2 a - \cos a$ , express  $f(a)$  in the form  $p \cos^2 a + q \cos a + r$ .

Hence solve, correct to three decimal places, the equation  $6 \sin^2 a - \cos a = 5$  for  $0 \leq a \leq \pi$ . 4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional		2.3 Source 1993 P1 qu.17
		C	A/B	C	A/B	C	A/B	2.3.1		
4	2.3			2	2					

$\bullet^1$  subst. leading from  $\sin^2$  to  $\cos^2$   
 $\bullet^2 -6\cos^2 a - \cos a + 6 = 5$   
 $\bullet^3$  solving the quadratic  
 $\bullet^4 1.231$  and  $2.094$

[SQA] 12. Find the values of  $t$ , where  $0 < t < 2\pi$ , for which  $4 \cos(2t - \frac{\pi}{4})$  has its maximum value. 4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional		2.3 Source 1989 P1 qu.15
		C	A/B	C	A/B	C	A/B	2.3.1		
4	2.3	4								

$\bullet^1 \cos(2t - \frac{\pi}{4}) = 1$   
 $\bullet^2 2t - \frac{\pi}{4} = 0$   
 $\bullet^3 t = \frac{\pi}{8}$   
 $\bullet^4 \frac{\pi}{8}, \frac{9\pi}{8}$

[SQA] 13. Solve the equation  $2 \sin\left(2x - \frac{\pi}{6}\right) = 1$ ,  $0 \leq x < 2\pi$ .

4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1998 P1 qu.9
		C	A/B	C	A/B	C	A/B		
.	4	4						2.3.1 1.2.1	


- <sup>1</sup>  $\sin\left(2x - \frac{\pi}{6}\right) = \frac{1}{2}$  Alternative for 2nd and 3rd marks  
 •<sup>2</sup>  $2x - \frac{\pi}{6} = \frac{\pi}{6}, \frac{5\pi}{6}$  (accept 30, 150)  
 •<sup>3</sup>  $x = \frac{\pi}{6}, \frac{\pi}{2}$   
 •<sup>4</sup>  $x = \frac{7\pi}{6}, \frac{3\pi}{2}$

[SQA] 14. (a) Solve the equation  $\sin 2x^\circ - \cos x^\circ = 0$  in the interval  $0 \leq x \leq 180$ .

4

(b) The diagram shows parts of two trigonometric graphs,  $y = \sin 2x^\circ$  and  $y = \cos x^\circ$ .

Use your solutions in (a) to write down the coordinates of the point P.




1

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
(a)	4	C	NC	T10	30, 90, 150	2001 P1 Q5
(b)	1	C	NC	T3	$(150, -\frac{\sqrt{3}}{2})$	

<ul style="list-style-type: none"> <li>•<sup>1</sup> ss: use double angle formula</li> <li>•<sup>2</sup> pd: factorise</li> <li>•<sup>3</sup> pd: process</li> <li>•<sup>4</sup> pd: process</li> <li>•<sup>5</sup> ic: interpret graph</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>2 \sin x^\circ \cos x^\circ</math></li> <li>•<sup>2</sup> <math>\cos x^\circ(2 \sin x^\circ - 1)</math></li> <li>•<sup>3</sup> <math>\cos x^\circ = 0, \sin x^\circ = \frac{1}{2}</math></li> <li>•<sup>4</sup> 90, 30, 150</li> </ul> <p>or</p> <ul style="list-style-type: none"> <li>•<sup>3</sup> <math>\sin x^\circ = \frac{1}{2}</math> and <math>x = 30, 150</math></li> <li>•<sup>4</sup> <math>\cos x^\circ = 0</math> and <math>x = 90</math></li> <li>•<sup>5</sup> <math>(150, -\frac{\sqrt{3}}{2})</math></li> </ul>
--	--

- [SQA] 15. The diagram shows two curves with equations  $y = \cos 2x^\circ$  and  $y = 1 + \sin x^\circ$  where  $0^\circ \leq x \leq 360^\circ$ .

Find the  $x$ -coordinate of the point of intersection at A.



4

part	marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3 Source 1991 P1 qu.20
			C	A/B	C	A/B	C	A/B		
.	4	2.3	1	3					2.3.5	

- <sup>1</sup>  $\cos 2x^\circ = 1 + \sin x^\circ$
- <sup>2</sup>  $2\sin^2 x^\circ + \sin x^\circ = 0$
- <sup>3</sup>  $\sin x^\circ = 0 \text{ or } -\frac{1}{2}$
- <sup>4</sup>  $x = 210$

- [SQA] 16. Functions  $f$  and  $g$  are defined on suitable domains by  $f(x) = \sin(x^\circ)$  and  $g(x) = 2x$ .

(a) Find expressions for:

- (i)  $f(g(x))$ ;
- (ii)  $g(f(x))$ .

(b) Solve  $2f(g(x)) = g(f(x))$  for  $0^\circ \leq x \leq 360^\circ$ .

2

5

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
(a)	2	C	CN	A4	(i) $\sin(2x^\circ)$ , (ii) $2\sin(x^\circ)$	2002 P1 Q3
(b)	5	C	CN	T10	$0^\circ, 60^\circ, 180^\circ, 300^\circ, 360^\circ$	

- <sup>1</sup> ic: interpret  $f(g(x))$
- <sup>2</sup> ic: interpret  $g(f(x))$
- <sup>3</sup> ss: equate for intersection
- <sup>4</sup> ss: substitute for  $\sin 2x$
- <sup>5</sup> pd: extract a common factor
- <sup>6</sup> pd: solve a 'common factor' equation
- <sup>7</sup> pd: solve a 'linear' equation


- <sup>1</sup>  $\sin(2x^\circ)$
- <sup>2</sup>  $2\sin(x^\circ)$
- <sup>3</sup>  $2\sin(2x^\circ) = 2\sin(x^\circ)$
- <sup>4</sup> appearance of  $2\sin(x^\circ)\cos(x^\circ)$
- <sup>5</sup>  $2\sin(x^\circ)(2\cos(x^\circ) - 1)$
- <sup>6</sup>  $\sin(x^\circ) = 0 \text{ and } 0, 180, 360$
- <sup>7</sup>  $\cos(x^\circ) = \frac{1}{2} \text{ and } 60, 300$

or

- <sup>6</sup>  $\sin(x^\circ) = 0 \text{ and } \cos(x^\circ) = \frac{1}{2}$
- <sup>7</sup>  $0, 60, 180, 300, 360$

[SQA] 17. (a) Solve the equation  $3\sin 2x^\circ = 2\sin x^\circ$  for  $0 \leq x \leq 360$  (4)

- (b) The diagram below shows parts of the graphs of sine functions  $f$  and  $g$ . State expressions for  $f(x)$  and  $g(x)$ . (1)
- (c) Use your answers to part (a) to find the co-ordinates of A and B. (2)
- (d) Hence state the values of  $x$  in the interval  $0 \leq x \leq 360$  for which  $3\sin 2x^\circ < 2\sin x^\circ$ . (3)



part	marks	Unit	non-calc		calc		calc neut		Content Reference :		2.3
			C	A/B	C	A/B	C	A/B	Main	Additional	
(a)	4	2.3			4				2.3.5		
(b)	1	1.2			1				1.2.7		
(c)	2	1.2			2				1.2.9		
(d)	3	1.2			2	1			1.2.10		
										Source 1992 Paper 2 Qu.7	


- (a) •<sup>1</sup> strategy: ie  $\sin 2x = 2\sin x \cos x$   
•<sup>2</sup>  $\sin x = 0$  AND  $\cos x = \frac{1}{3}$   
•<sup>3</sup> 0, 180 AND 360  
•<sup>4</sup> 70.5 AND 289.5 AND no other angles
- (b) •<sup>5</sup>  $f(x) = 2\sin x^\circ$ ,  $g(x) = 3\sin 2x^\circ$
- (c) •<sup>6</sup>  $x = 70.5$  AND  $289.5$   
•<sup>7</sup>  $y = 1.89$  AND  $-1.89$
- (d) •<sup>8</sup> 70.5 AND 180  
•<sup>9</sup> 289.5 AND 360  
•<sup>10</sup> use inequality signs logically to connect the points of intersection (ie not for  $180 < x < 70.5$ )

- [SQA] 18. The diagram shows the graph of a cosine function from 0 to  $\pi$ .

(a) State the equation of the graph.

(b) The line with equation  $y = -\sqrt{3}$  intersects this graph at point A and B.

Find the coordinates of B.



1


3

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
(a)	1	C	NC	T4	$y = 2 \cos 2x$	2002 P1 Q8
(b)	3	C	NC	T7	$B(\frac{7\pi}{12}, -\sqrt{3})$	

- <sup>1</sup> ic: interpret graph
- <sup>2</sup> ss: equate equal parts
- <sup>3</sup> pd: solve linear trig equation in radians
- <sup>4</sup> ic: interpret result

- <sup>1</sup>  $2 \cos 2x$
- <sup>1</sup>  $2 \cos 2x = -\sqrt{3}$
- <sup>2</sup>  $2x = \frac{5\pi}{6}, \frac{7\pi}{6}$
- <sup>3</sup>  $x = \frac{7\pi}{12}$

- [SQA] 19. The diagram shows part of the graph of  $y = a \sin 3x^\circ$  and the line with equation  $y = \frac{2}{3}a$ . Find the x-coordinates of A and B.



4

part	marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1999 P1 qu.14
			C	A/B	C	A/B	C	A/B		
.	4	2.3	4						2.3.1	


- <sup>1</sup>  $a \sin 3x = \frac{2}{3}a$  stated or implied by •<sup>2</sup>
- <sup>2</sup>  $\sin 3x = \frac{2}{3}$
- <sup>3</sup>  $3x = 41.8, 138.2$  (138.2 stated or implied by 46.1 in •<sup>4</sup>)
- <sup>4</sup> 13.9, 46.1

- [SQA] 20. The diagram shows the graph of a sine function from  $0^\circ$  to  $90^\circ$ .

(a) State the equation of the graph.

(b) The line with equation  $y = -1.5$  intersects the curve at A and B.

Find the coordinates of A and B.



2


3

part	marks	Unit	non-calc		calc		calc neut		Content Reference :		
			C	A/B	C	A/B	C	A/B	Main	Additional	
(a)	2	1.2			2				1.2.2	1.2.7	
(b)	3	2.3			3				2.3.1		Source 1990 P1 qu.10

- <sup>1</sup>  $\sin 4x$
- <sup>2</sup> (trig function)  $\times 2$
- <sup>3</sup>  $f(x) = -1.5$
- <sup>4</sup>  $57.1^\circ$
- <sup>5</sup>  $77.9^\circ$

07

- [SQA] 21. The sketch represents part of the graph of a trigonometric function of the form  $y = p \sin(x + r)^\circ + q$ . It crosses the axes at  $(0, s)$  and  $(t, 0)$ , and has turning points at  $(50, -2)$  and  $(u, 4)$ .
- Write down values for  $p$ ,  $q$ ,  $r$  and  $u$ . (4)
  - Find the values for  $s$  and  $t$ . (4)




part	marks	Unit	non-calc		calc		calc neut		Content Reference :		2.3
			C	A/B	C	A/B	C	A/B	Main	Additional	
(a)	4	1.2			2	2			1.2.3		
(b)	4	2.3				4			2.3.1		Source 1997 Paper 2 Qu.9

(a) •<sup>1</sup>  $p = -3$   
•<sup>2</sup>  $q = 1$   
•<sup>3</sup>  $r = 40$  or  $-320$   
•<sup>4</sup>  $u = 230$

(b) •<sup>5</sup> replace  $x$  by 0  
•<sup>6</sup>  $-0.928$   
•<sup>7</sup> replace  $y$  by 0  
•<sup>8</sup>  $120.5$

- [SQA] 22. The diagram shows the graph of the function  $y = a + b \sin cx$  for  $0 \leq x \leq \pi$ .

- (a) Write down the values of  $a$ ,  $b$  and  $c$ .  
 (b) Find algebraically the values of  $x$  for which  $y = 2.5$ .

3  
3


part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
(a) 3	1.2	3						1.2.3	
(b) 3	2.3	3						2.3.1 1.2.11	1994 P1 qu.12

$$\begin{array}{ll} \bullet^1 a = 2 & \bullet^4 2 + \sin 2x = 2\frac{1}{2} \\ \bullet^2 b = 1 & \bullet^5 2x = \frac{\pi}{6}, \frac{5\pi}{6} \\ \bullet^3 c = 2 & \bullet^6 x = \frac{\pi}{12}, \frac{5\pi}{12} (0.262, 1.309) \end{array}$$

$$\text{OR} \quad \begin{array}{ll} \bullet^4 2 + \sin 2x = 2\frac{1}{2} \\ \bullet^5 2x = \frac{\pi}{6}, x = \frac{\pi}{12} \\ \bullet^6 2x = \frac{5\pi}{6}, x = \frac{5\pi}{12} \end{array}$$

- [SQA] 23. The diagram shows a sketch of the graph of  $y = \sin(2x - \frac{\pi}{6})$ ,  $0 \leq x \leq \pi$ , and the straight line  $y = 0.5$ . These graphs intersect at P and Q.

Find algebraically the coordinates of P and Q.




4

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
4	2.3	4						2.3.1 1.2.1	1996 P1 qu.12

$$\begin{array}{ll} \bullet^1 \sin(2x - \frac{\pi}{6}) = 0.5 & \text{stated or implied by 2nd mark} \\ \bullet^2 2x - \frac{\pi}{6} = \frac{\pi}{6} \\ \bullet^3 2x - \frac{\pi}{6} = \frac{5\pi}{6} \\ \bullet^4 (\frac{\pi}{6}, 0.5), (\frac{\pi}{2}, 0.5) \end{array}$$

- [SQA] 24. The diagram below shows the graph of  $y = 2\sin 2x + 1$  for  $0 \leq x \leq \pi$ .




- (a) Find the coordinates of A and B (as shown in the diagram) by solving an appropriate equation algebraically. (5)
- (b) The points  $(0, 2)$  and  $(\pi, 0)$  are joined by a straight line  $l$ . In how many points does  $l$  intersect the given graph? (1)
- (c) C is the point on the given graph with an x-coordinate of  $\frac{\pi}{2}$ . Explain whether C is above, below or on the line  $l$ . (3)

part	marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
			C	A/B	C	A/B	C	A/B		
(a)	5	2.3	3	2					2.3.1	
(b)	1	0.1	1						0.1	
(c)	3	0.1		3					0.1	Source 1993 Paper 2 Qu.6

- (a) •<sup>1</sup>  $2\sin 2x + 1 = 0$   
     •<sup>2</sup>  $\sin 2x = -\frac{1}{2}$   
     •<sup>3</sup> for any valid sol of equ. in form  $\sin ax = -\frac{b}{c}$   
     •<sup>4</sup>  $(\frac{7\pi}{12}, 0)$   
     •<sup>5</sup>  $(\frac{11\pi}{12}, 0)$
- (b) •<sup>6</sup> 3
- (c) •<sup>7</sup>  $y_C = 1$   
     •<sup>8</sup> for a strategy to make a decision about C  
     •<sup>9</sup> for making a consistent decision about C

[SQA] 25. Using triangle PQR, as shown, find the exact value of  $\cos 2x$ .

3



part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
.	3	2.3	3					2.3.3	Source 1999 P1 qu.12

•<sup>1</sup>  $\cos x = \frac{2}{\sqrt{11}}$  or  $\sin x = \frac{\sqrt{7}}{\sqrt{11}}$

•<sup>2</sup>  $\cos 2x = 2 \times \left(\frac{2}{\sqrt{11}}\right)^2 - 1$

•<sup>3</sup>  $-\frac{3}{11}$

[SQA] 26. If  $\cos \theta = \frac{4}{5}$ ,  $0 \leq \theta < \frac{\pi}{2}$ , find the exact value of

(a)  $\sin 2\theta$

2

(b)  $\sin 4\theta$ .

3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
(a)	2	2.3	2					2.3.3	Source
(b)	3	2.3		3				2.3.3	1994 P1 qu.13

•<sup>1</sup>  $\sin \theta = \frac{3}{5}$

•<sup>3</sup>  $2 \sin 2\theta \cos 2\theta$

•<sup>2</sup>  $\frac{24}{25}$

•<sup>4</sup>  $\cos 2\theta = \frac{7}{25}$

•<sup>5</sup>  $\frac{336}{625}$

[SQA] 27. Given that  $\tan \alpha = \frac{\sqrt{11}}{3}$ ,  $0 < \alpha < \frac{\pi}{2}$ , find the exact value of  $\sin 2\alpha$ .

3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
.	3	2.3	3					2.3.3	Source 1995 P1 qu.12

•<sup>1</sup> "third side" =  $\sqrt{20}$

•<sup>2</sup>  $\sin \alpha = \frac{\sqrt{11}}{\sqrt{20}}$  or  $\cos \alpha = \frac{3}{\sqrt{20}}$

•<sup>3</sup>  $2 \times \frac{\sqrt{11}}{\sqrt{20}} \times \frac{3}{\sqrt{20}}$

[SQA] 28. Given that  $\cos D = \frac{2}{\sqrt{5}}$  and  $0 < D < \frac{\pi}{2}$ , find the exact values of  $\sin D$  and  $\cos 2D$ . 3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1990 P1 qu.9
		C	A/B	C	A/B	C	A/B		
3	2.3	3						2.3.3	

- <sup>1</sup> strat for exact value: e.g.  $\sin^2 D = 1 - \cos^2 D$
- <sup>2</sup>  $\sin D = \frac{1}{\sqrt{5}}$
- <sup>3</sup>  $\cos 2D = \frac{3}{5}$

[SQA] 29. Given that  $\sin A = \frac{3}{4}$ , where  $0 < A < \frac{\pi}{2}$ , find the exact value of  $\sin 2A$ . 3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1991 P1 qu.12
		C	A/B	C	A/B	C	A/B		
3	2.3	3						2.3.3	

- <sup>1</sup> strat for cos: eg  $\cos^2 = 1 - \sin^2$
- <sup>2</sup>  $\cos A = \frac{\sqrt{7}}{4}$
- <sup>3</sup>  $\sin 2A = \frac{3\sqrt{7}}{8}$

[SQA] 30. For acute angles  $P$  and  $Q$ ,  $\sin P = \frac{12}{13}$  and  $\sin Q = \frac{3}{5}$ .

Show that the exact value of  $\sin(P + Q)$  is  $\frac{63}{65}$ . 3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1993 P1 qu.6
		C	A/B	C	A/B	C	A/B		
3	2.3	3						2.3.2	

- <sup>1</sup>  $\cos P = \frac{5}{13}$
- <sup>2</sup>  $\cos Q = \frac{4}{5}$
- <sup>3</sup>  $\frac{12}{13} \times \frac{4}{5} + \frac{5}{13} \times \frac{3}{5}$

[SQA] 31. Find the exact value of  $\sin \theta^\circ + \sin(\theta^\circ + 120^\circ) + \cos(\theta^\circ + 150^\circ)$ .

3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1994 P1 qu.6
		C	A/B	C	A/B	C	A/B		
3	2.3	3						2.3.2	

- <sup>1</sup>  $\sin \theta \cos 120 + \cos \theta \sin 120$  and  $\cos \theta \cos 150 - \sin \theta \sin 150$
- <sup>2</sup> correct use of exact values
- <sup>3</sup> simplification to zero

[SQA] 32. If  $x^\circ$  is an acute angle such that  $\tan x^\circ = \frac{4}{3}$ , show that the exact value of  $\sin(x^\circ + 30^\circ)$  is  $\frac{4\sqrt{3}+3}{10}$ .


3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1997 P1 qu.7
		C	A/B	C	A/B	C	A/B		
3	2.3	3						2.3.2	

- <sup>1</sup>  $\sin x^\circ \cos 30^\circ + \cos x^\circ \sin 30^\circ$
- <sup>2</sup>  $\sin x^\circ = \frac{4}{5}$  &  $\cos x^\circ = \frac{3}{5}$
- <sup>3</sup>  $\frac{4}{5} \cdot \frac{\sqrt{3}}{2} + \frac{3}{5} \cdot \frac{1}{2}$  and completes proof

- [SQA] 33. The diagram shows two right-angled triangles ABD and BCD with AB = 7 cm, BC = 4 cm and CD = 3 cm. Angle DBC =  $x^\circ$  and angle ABD =  $y^\circ$ .

Show that the exact value of  $\cos(x + y)^\circ$  is  $\frac{20 - 6\sqrt{6}}{35}$ .



3

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
. 3	2.3					3		2.3.2	Source 1996 P1 qu.15

- <sup>1</sup> know to calculate missing sides
- <sup>2</sup>  $BD = 5$ ,  $AD = \sqrt{24}$
- <sup>3</sup>  $\cos x \cos y - \sin x \sin y = \frac{4}{5} \cdot \frac{5}{7} - \frac{3}{5} \cdot \frac{\sqrt{24}}{7}$

- [SQA] 34.  $A$  and  $B$  are acute angles such that  $\tan A = \frac{3}{4}$  and  $\tan B = \frac{5}{12}$ .

Find the exact value of

(a)  $\sin 2A$

2

(b)  $\cos 2A$

1

(c)  $\sin(2A + B)$ .

2

part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3
		C	A/B	C	A/B	C	A/B		
(a) 2	2.3	2						2.3.3	
(b) 1	2.3	1						2.3.3	
(c) 2	2.3	2						2.3.2	Source 1998 P1 qu.7

•<sup>1</sup>  $\sin A = \frac{3}{5}$  and  $\cos A = \frac{4}{5}$

•<sup>4</sup>  $\sin 2A \cos B + \cos 2A \sin B$

•<sup>2</sup>  $\sin 2A = 2 \times \frac{3}{5} \times \frac{4}{5} = \frac{24}{25}$  (accept 0.96)

•<sup>5</sup>  $\sin B = \frac{5}{13}$  and  $\cos B = \frac{12}{13}$  and  $\frac{123}{325}$

•<sup>3</sup>  $\cos 2A = \text{e.g. } \left(\frac{4}{5}\right)^2 - \left(\frac{3}{5}\right)^2 = \frac{7}{25}$  (accept 0.28)

- [SQA] 35. Functions  $f(x) = \sin x$ ,  $g(x) = \cos x$  and  $h(x) = x + \frac{\pi}{4}$  are defined on a suitable set of real numbers.

(a) Find expressions for:

(i)  $f(h(x))$ ;

(ii)  $g(h(x))$ .

2

(b) (i) Show that  $f(h(x)) = \frac{1}{\sqrt{2}} \sin x + \frac{1}{\sqrt{2}} \cos x$ .

(ii) Find a similar expression for  $g(h(x))$  and hence solve the equation  
 $f(h(x)) - g(h(x)) = 1$  for  $0 \leq x \leq 2\pi$ .


5

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
(a)	2	C	NC	A4	(i) $\sin(x + \frac{\pi}{4})$ , (ii) $\cos(x + \frac{\pi}{4})$	2001 P1 Q7
(b)	5	C	NC	T8, T7	(i) proof, (ii) $x = \frac{\pi}{4}, \frac{3\pi}{4}$	

<ul style="list-style-type: none"> <li>•<sup>1</sup> ic: interpret composite functions</li> <li>•<sup>2</sup> ic: interpret composite functions</li> <li>•<sup>3</sup> ss: expand <math>\sin(x + \frac{\pi}{4})</math></li> <li>•<sup>4</sup> ic: interpret</li> <li>•<sup>5</sup> ic: substitute</li> <li>•<sup>6</sup> pd: start solving process</li> <li>•<sup>7</sup> pd: process</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>\sin(x + \frac{\pi}{4})</math></li> <li>•<sup>2</sup> <math>\cos(x + \frac{\pi}{4})</math></li> <li>•<sup>3</sup> <math>\sin x \cos \frac{\pi}{4} + \cos x \sin \frac{\pi}{4}</math> and complete</li> <li>•<sup>4</sup> <math>g(h(x)) = \frac{1}{\sqrt{2}} \cos x - \frac{1}{\sqrt{2}} \sin x</math></li> <li>•<sup>5</sup> <math>(\frac{1}{\sqrt{2}} \sin x + \frac{1}{\sqrt{2}} \cos x) - (\frac{1}{\sqrt{2}} \cos x - \frac{1}{\sqrt{2}} \sin x)</math></li> <li>•<sup>6</sup> <math>\frac{2}{\sqrt{2}} \sin x</math></li> <li>•<sup>7</sup> <math>x = \frac{\pi}{4}, \frac{3\pi}{4}</math> accept only radians</li> </ul>
--	---

- [SQA] 36. On the coordinate diagram shown, A is the point  $(6, 8)$  and B is the point  $(12, -5)$ . Angle  $AOC = p$  and angle  $COB = q$ .

Find the exact value of  $\sin(p + q)$ .




4

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
	4	C	NC	T9	$\frac{63}{65}$	2000 P1 Q1

- <sup>1</sup> ss: know to use trig expansion
- <sup>2</sup> pd: process missing sides
- <sup>3</sup> ic: interpret data
- <sup>4</sup> pd: process

- <sup>1</sup>  $\sin p \cos q + \cos p \sin q$
- <sup>2</sup> 10 and 13
- <sup>3</sup>  $\frac{8}{10} \cdot \frac{12}{13} + \frac{6}{10} \cdot \frac{5}{13}$
- <sup>4</sup>  $\frac{126}{130}$

- [SQA] 37. In triangle ABC, show that the exact value of  $\sin(a + b)$  is  $\frac{2}{\sqrt{5}}$ .



4

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
	4	C	NC	T9	proof	2002 P1 Q5


- <sup>1</sup> pd: process the missing sides
- <sup>2</sup> ss: expand
- <sup>3</sup> pd: substitute
- <sup>4</sup> pc: process and complete proof

- <sup>1</sup>  $AC = \sqrt{2}$  and  $BC = \sqrt{10}$  stated or implied by •<sup>3</sup>
- <sup>2</sup>  $\sin(a + b) = \sin a \cos b + \cos a \sin b$
- <sup>3</sup>  $\frac{1}{\sqrt{2}} \cdot \frac{3}{\sqrt{10}} + \frac{1}{\sqrt{2}} \cdot \frac{1}{\sqrt{10}}$
- <sup>4</sup>  $\frac{4}{\sqrt{20}} = \dots = \frac{2}{\sqrt{5}}$

[SQA] 38. The diagram shows an isosceles triangle PQR in which PR = QR and angle PQR =  $x^\circ$ .

(a) Show that  $\frac{\sin x^\circ}{p} = \frac{\sin 2x^\circ}{r}$ .

- (b) (i) State the value of  $x^\circ$  when  $p = r$ .  
(ii) Using the fact that  $p = r$ , solve the equation in (a) above, to justify your stated value of  $x^\circ$ .




part	marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1991 Paper 2 Qu. 3
			C	A/B	C	A/B	C	A/B		
(a)	3	0.1					2	1	0.1	
(b)	5	2.3					5		2.3.5, 0.1	

(a) •<sup>1</sup>  $(180 - 2x)^\circ$   
•<sup>2</sup>  $\frac{\sin x^\circ}{p} = \frac{\sin(180 - 2x)^\circ}{r}$   
•<sup>3</sup>  $\sin(180 - 2x)^\circ = \sin 2x^\circ$  stated explicitly

(b) •<sup>4</sup>  $60^\circ$   
•<sup>5</sup>  $\sin x^\circ = \sin 2x^\circ$   
•<sup>6</sup>  $\sin x^\circ(2 \cos x^\circ - 1) = 0$   
•<sup>7</sup>  $\sin x^\circ = 0$  and  $\cos x^\circ = \frac{1}{2}$   
•<sup>8</sup>  $x = 60$  is only answer stated explicitly

- [SQA] 39. The framework of a child's swing has dimensions as shown in the diagram on the right.  
Find the exact value of  $\sin x^\circ$ .

5



part marks	Unit	non-calc		calc		calc neut		Content Reference : Main Additional	2.3  Source 1996 P1 qu.18
		C	A/B	C	A/B	C	A/B		
. 5	2.3	1	4					2.3.4	

• <sup>1</sup>	sketch with $\frac{x}{2}$ marked in r/a $\Delta$	OR	• <sup>1</sup>	know to use cosine rule
• <sup>2</sup>	height of triangle = $\sqrt{5}$		• <sup>2</sup>	$\cos x = \frac{3^2 + 3^2 - 4^2}{2 \cdot 3 \cdot 3}$
• <sup>3</sup>	$\sin x = 2 \sin \frac{1}{2}x \cos \frac{1}{2}x$		• <sup>3</sup>	$\frac{1}{9}$
• <sup>4</sup>	$\sin \frac{x}{2} = \frac{2}{3}$ and $\cos \frac{1}{2}x = \frac{\sqrt{5}}{3}$		• <sup>4</sup>	draw r/a $\Delta$ or use $\cos^2 x + \sin^2 x = 1$
• <sup>5</sup>	$\sin x = \frac{4\sqrt{5}}{9}$		• <sup>5</sup>	$\sin x = \frac{\sqrt{80}}{9}$

[END OF WRITTEN QUESTIONS]